

# Cyanocobalamin (Vitamin B12) 1000 µg/ml Solution for Injection

Cyanocobalamin (Vitamin B12) 1000 micrograms in 1 ml. Phenol 0.5% w/v (as a preservative).

**Uses:** For intramuscular or subcutaneous injection. As an aid in raising blood Vitamin B12 levels in foals, calves, horses and cattle. For use in the treatment of Vitamin B12 deficiency, and for poor growth rates and general unthriftiness in young animals when associated with the above deficiency.

**Dosage:** Foals and Calves: 0.5-1 ml. Horses and Cattle: 1-3 ml. Repeat in 7 days if required.

**Warnings:** Do not give by the intravenous route. Observe normal aseptic precautions. Following withdrawal of the first dose, use the product within 28 days. Do not store above 25°C. Protect from light. Keep out of the reach and sight of children. For animal treatment only.

**Withdrawal period:** Cattle, horse meat: zero days.

Cattle milk: zero hours.

Animalcare Ltd, York, YO26 6RB, UK

Vm 10434/4064 E1747

[POM-VPS] GTIN: 05055037400106

---

## Presentation:

A clear, red, sterile solution for injection containing Cyanocobalamin (Vitamin B12) 1000 micrograms in 1 ml. The product contains Phenol 0.5% w/v as a preservative.

## Uses:

As an aid in raising blood Vitamin B12 levels in horses and cattle.

For use in the treatment of Vitamin B12 deficiency, and for poor growth rates and general unthriftiness in young animals when associated with the above deficiency.

## DOSAGE AND ADMINISTRATION

**Calves and foals: 0.5-1 ml    Horses and cattle: 1-3 ml**

By intramuscular or subcutaneous injection. Repeat in 7 days if required.

---

## Contraindications:

Do not give by the intravenous route.

## General precautions:

Care should be taken to avoid accidental self-injection. If accidental self-injection occurs, seek medical advice immediately. In case of accidental ingestion, seek medical advice immediately. In case of accidental spillage onto the skin or eyes, wash the affected area immediately with clean running water. Seek medical attention if irritation persists.

Keep out of the reach and sight of children. For animal treatment only.

Observe normal aseptic precautions.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

---

---

**Withdrawal period:**

Cattle and horse meat: zero days.

Cattle milk: zero hours.

**Pharmaceutical precautions:**

Do not store above 25 °C. Protect from light.

Following withdrawal of the first dose, use the product within 28 days.

Rinse containers thoroughly with water. Dispose of rinsings in slurry or dirty water. Dispose of rinsed containers in the farm refuse. Used containers should not be recycled. When the container is broached for the first time, using the in-use shelf-life which is specified on this package leaflet, the date on which any product remaining in the container should be discarded should be worked out. This discard date should be written in the space provided on the label.

---

**Legal category:**

**POM-VPS** Prescription Only Medicine - Veterinarian, Pharmacist, Suitably Qualified Person

To be supplied only on veterinary prescription.

UK authorised veterinary medicinal product.

**Package quantity:**

Multidose vial of 50 ml.

**Further information:**

Following treatment, the Vitamin B12 status of herbivores should be maintained by dietary supplementation with Cobalt.

---

**Marketing authorisation number:**

Vm 10434/4064

**Date of last approval:**

January 2014

Animalcare Ltd,  
10 Great North Way,  
York,  
YO26 6RB  
UK