

Anivit 4BC Solution for Injection

STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS

Anivit 4BC is a pale yellow to brown sterile solution for injection containing four members of the B-complex vitamins together with vitamin C.

Thiamine hydrochloride (Vitamin B1)	3.5% w/v
Riboflavine sodium phosphate (Vitamin B2)	0.05% w/v
Pyridoxine hydrochloride (Vitamin B6)	0.7% w/v
Nicotinamide	2.3% w/v
Ascorbic acid (Vitamin C)	7.0% w/v
Chlorocresol (preservative)	0.1% w/v

INDICATION(S)

Anivit 4BC may be used in the following cases:

- 1) Wherever deficiency of vitamin B group is suspected in horses, cattle and sheep.
- 2) In treatment of cerebrocortical necrosis in cattle and sheep.
- 3) In treatment of bracken poisoning in horses.

CONTRAINDICATIONS

Anaphylactoid reactions, particularly in the horse may occur following intravenous administration. When this route is used Anivit 4BC should be given slowly and may be diluted with sterile saline or sterile dextrose saline solution. This may be achieved by drawing sterile diluting solution into a syringe, followed by the required volume of Anivit 4BC. The diluted product must be administered immediately after preparation and must not be stored. Aseptic precautions should be observed. There may be slight irritation at the site of injection when the product is given by the subcutaneous or intramuscular routes.

ADVERSE REACTIONS

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

TARGET SPECIES

Horses
Cattle
Sheep

DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Administer by subcutaneous, deep intramuscular or slow intravenous injection. The dose should be repeated daily as required.

SPECIES	DOSAGE
Horses, cattle	20 - 30 ml
Calves, foals	5 - 10 ml
Sheep	5 - 10 ml

ADVICE ON CORRECT ADMINISTRATION

Normal aseptic precautions should be observed.
If dose volume exceeds 20ml, it should be divided into two sites.

WITHDRAWAL PERIOD

Withdrawal periods: Meat: zero days; Milk: zero hours

USER WARNINGS

Care should be taken to avoid accidental self-injection.
Wash hands after use.

SPECIAL STORAGE PRECAUTIONS

Do not store above 25°C.

Protect from light.

Keep out of the reach and sight of children.

Keep container in outer carton.

Following withdrawal of the first dose, use the product within 28 days. Discard unused material.

When the container is breached (opened) for the first time, using the in-use shelf-life which is specified on this package leaflet, the date on which any product remaining in the carton should be discarded should be worked out. This discard date should be written in the space provided.

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

October 2008

OTHER INFORMATION

Multidose vials of 50ml and 100 ml.

POM-VPS

Vm 02000/4219

To be supplied only on veterinary prescription

NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Norbrook Laboratories Limited,
Station Works, Newry,
Co Down, BT35 6JP,
Northern Ireland

DISTRIBUTED BY:

Animalcare Ltd.
10 Great North Way
York, YO26 6RB

FOR ANIMAL TREATMENT ONLY