

oxycare[®] 50mg

Oxycare[®] For the treatment of oxytetracycline sensitive bacterial disease in Dogs
Oxytetracycline dihydrate 50mg

Each tablet contains as active ingredient:
Oxytetracycline dihydrate 50mg.

Use: For the treatment of oxytetracycline sensitive bacterial disease.

Dosage and administration: In dogs: Initial dose of 50mg/kg body weight followed by 25mg/kg every 12 hours for 5 days. For oral administration at least one hour before or two hours after feeding.

Contra-indications/warnings: Contraindicated in patients with hypersensitivity to any tetracyclines. Do not administer concurrently with milk or antacids. Oxytetracycline is deposited in growing teeth and bones and may cause yellow discolouration. It also crosses the placenta. For this reason it is not recommended in late pregnancy or in young animals. Caution must be taken in treating animals with renal or hepatic dysfunction; in such cases it may be necessary to reduce dosage levels. If you know you are hypersensitive (allergic) to oxytetracycline, do not handle the product. In the event of accidental ingestion, flush mouth with plenty of water and seek medical advice. In the event of eye contact, flush thoroughly with clean, running water. If irritation persists seek medical attention. Wash hands after use.

Disposal advice: Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

Store below 25°C in a dry place.

Protect from light.

Keep out of the reach and sight of children.

For animal treatment only.

To be supplied only on veterinary prescription.

Legal category: POM-V

Product license number: Vm 10347/4001. E1931.

Marketing authorisation holder:

Animalcare Ltd, 10 Great North Way, York, YO26 6RB, UK

UK only:

Vm 10347/4001

POM-V Prescription only medicine